

<b>SUSPECTED CVA/TIA</b>	
<b>ADULT</b>	<b>PEDIATRIC (≤34KG)</b>
<b>BLS</b>	
<ul style="list-style-type: none"> <li>• Universal Protocol #601</li> <li>• Pulse Oximetry                             <ul style="list-style-type: none"> <li>○ O<sub>2</sub> administration per Airway Management Protocol #602</li> </ul> </li> <li>• Place patient in position of comfort with head elevated</li> <li>• Complete B-E-F-A-S-T exam, see Notes</li> </ul>	Same as Adult
<b>BLS Elective Skills</b>	
Obtain Blood Sugar Level – if <60 mg/dL see Altered Mental Status Protocol #612	
<b>ALS Standing Orders</b>	
<ul style="list-style-type: none"> <li>• Initiate <b>STROKE ALERT</b> base report and EARLY transport if ANY of B-E-F-A-S-T present <b>and</b> last seen normal is &lt; 6 hours</li> </ul>	Same as Adult
<b>Base Hospital Orders Only</b>	
<ul style="list-style-type: none"> <li>• As needed</li> </ul>	<ul style="list-style-type: none"> <li>• As needed</li> </ul>
<b>Notes</b>	
<ul style="list-style-type: none"> <li>• Contact receiving hospital early if patient meets STROKE ALERT criteria</li> <li>• Do not delay transport to hospital for on scene treatment</li> <li>• B-E-F-A-S-T ( new onset or change from previous “normal”)(*)                             <ul style="list-style-type: none"> <li>○ B – Balance – changes or problems</li> <li>○ E – Eyes – sudden change in vision or double vision</li> <li>○ F – Facial droop</li> <li>○ A – Arm drift</li> <li>○ S – Speech abnormalities</li> <li>○ T – Time last seen normal &lt; 6 hrs. (record the time)</li> </ul> </li> <li>• “Time last seen normal” – do not confuse with time symptoms noticed – “time last seen normal” starts the clock</li> <li>• Evaluate and correct for other potential causes i.e.                             <ul style="list-style-type: none"> <li>○ Hypoxia</li> <li>○ Hypoglycemia</li> <li>○ Overdose</li> <li>○ Seizure</li> </ul> </li> <li>• If time is available and the patient or family can provide the information, gather the criteria listed in the Fibrinolytic Evaluation listed below (Fibrinolytic Evaluation – ASA/NSA Standards)                             <ul style="list-style-type: none"> <li>○ Use of anticoagulants</li> <li>○ History of stroke/TIA/brain tumor</li> <li>○ Mental Status - orientation</li> <li>○ Known bleeding disorder</li> <li>○ Age &gt; 18 yrs.</li> <li>○ Active bleeding, surgery, or trauma &lt; 3 weeks</li> <li>○ Jaundice, hepatitis, kidney failure</li> <li>○ Terminal illness</li> </ul> </li> </ul>	
<p>(*) BEFAST – was developed by Intermountain Healthcare, as an adaptation of the FAST model implemented by the American Stroke Assoc. Reproduction with the permission from Intermountain Healthcare, Copyright 2011, Intermountain Health Care.</p>	