County of San Luis Obispo Public Health Department Division: Emergency Medical Services Agency

**CHILDBIRTH** Normal Complications BLS Universal Protocol #601 Universal Protocol #601 Pulse Oximetry Pulse Oximetry O O2 administration per Airway O O2 administration per Airway Management Protocol #602 Management Protocol #602 Delivery Initiate Transport Early Patient assessment with visual exam of Hypertension BP > 180/110 mmHg perineum for crowning Seizures - follow Seizure Protocol #620 Control head and speed of delivery Check for cord around neck Vaginal bleeding in last trimester not Deliver upper, then lower shoulder associated with labor may indicate Placenta Dry, stimulate, and wrap baby Previa/Abruptio Suction airway as needed Breech/Limb presentation Cut and clamp cord 6" from newborn's Prolapsed cord - Place mother in knee-chest umbilicus position. Feel cord for pulse. With gloved Healthy infant to mother's breast hand, push baby into vagina slightly to take Prepare for delivery of placenta pressure off cord. Maintain this position. Do not attempt to push cord back Postpartum Hemorrhage Control Perform visual exam to determine site of bleeding For perineal tear, apply direct pressure Firmly massage fundus **ALS Standing Orders** Refer to appropriate protocol based on • Refer to appropriate protocol based on patient's presentation patient's presentation **Base Hospital Orders Only** As needed As needed **Notes** Knee-chest position for prolapse chord presentation

Protocol #650

Effective Date: 08/01/2019

- General guideline transport in left lateral position
- Obtain Para/Gravida