

Operations Subcommittee

of the Emergency Medical Care Committee



Meeting Agenda:
9 A.M., Thursday April 4th, 2024
Location: SLOEMSA Conference Room
2995 McMillan Ave, STE #178, San Luis Obispo

Members

Jay Wells, *Sheriff's Department, CHAIR*
 Tim Nurge, *Ambulance Providers*
 Scotty Jalbert, *Office of Emergency Services*
 Jennifer Mebane, *Med-Com*
 Adam Forrest, M.D., *Hospitals*
 Kris Strommen, *Ambulance Providers*
 Rob Jenkins, *Fire Service*
 Lisa Epps, *Air Ambulance Providers*
 Dennis Rowley, *Air Ambulance Providers*
 Doug Weeda, *CHP*
 Deputy Chief Sammy Fox, *Fire Service*
Vacant, Law Enforcement
 Chief Casey Bryson, *Fire Service*
 Chief Dan McCrain, *Fire Service*
 Roger Colombo, *Field Provider-Paramedic*

Staff

STAFF LIAISON, Ryan Rosander, *EMS Coordinator*
 Bill Mulkerin, M.D., *Medical Director*
 Rachel Oakley, *EMS Coordinator*
Vacant, EMS Coordinator
 Alyssa Vardas, *Administrative Assistant*

AGENDA	ITEM	LEAD
Call to Order	Introductions Public Comment	Jay Wells
Summary Notes	Review of Summary Notes February 1 st , 2024	
Discussion	<p>Protocol and Procedure Revisions:</p> <ul style="list-style-type: none"> Revised Protocol #602: Airway Management Revised Protocol #641: Cardiac Arrest Atraumatic Revised Protocol #661: Traumatic Cardiac Arrest Revised Procedure #717: Endotracheal Intubation Revised Procedure #718: Supraglottic Airway Device 	Ryan Rosander
Adjourn	<p>Declaration of Future Agenda Items -Roundtable</p> <hr/> <p>Next Meeting Date: TBD, 2024, 9:00 A.M. Location: SLOEMSA Conference Room 2995 McMillan Ave, STE #178, San Luis Obispo</p>	Jay Wells

DRAFT

Operations Subcommittee of the Emergency Medical Care Committee



Meeting Minutes

Thursday, February 1st, 2024

SLO EMSA Conference Room – 2995 McMillan Ave, Suite 178, San Luis Obispo

Members		Staff	
<input type="checkbox"/>	CHAIR Jay Wells, Sheriff's Department	<input checked="" type="checkbox"/>	STAFF LIASON Ryan Rosander, EMS Coordinator
<input type="checkbox"/>	Tim Benes, Ambulance Providers	<input checked="" type="checkbox"/>	Bill Mulkerin, MD, Medical Director
<input checked="" type="checkbox"/>	Scotty Jalbert, OES	<input checked="" type="checkbox"/>	Rachel Oakley, EMS Coordinator
<input checked="" type="checkbox"/>	Rob Jenkins, Fire Service	<input type="checkbox"/>	Vacant, EMS Coordinator
<input type="checkbox"/>	Adam Forrest, MD, Hospitals	<input checked="" type="checkbox"/>	Alyssa Vardas, EMS Administrative Assistant
<input checked="" type="checkbox"/>	Steve Lieberman, Fire Service		
<input checked="" type="checkbox"/>	Kris Strommen, Ambulance Providers		
<input type="checkbox"/>	Lisa Epps, Air Ambulance Providers		
<input checked="" type="checkbox"/>	Chief Casey Bryson, Fire Service		
<input type="checkbox"/>	Gerry Perez, CHP		
<input type="checkbox"/>	Chief Sammy Fox, Fire Service		
<input checked="" type="checkbox"/>	Roger Colombo, Field Provider, Paramedics		
<input checked="" type="checkbox"/>	Jessica Blaylock, Med-Com		
<input type="checkbox"/>	Aaron Hartney, Air Ambulance Providers		
<input type="checkbox"/>	Chief Casey Bryson, Fire Service		
<input checked="" type="checkbox"/>	Pete Gavitte, CHP		
<input type="checkbox"/>	Vacant, Law Enforcement		

AGENDA ITEM / DISCUSSION	ACTION / FOLLOW-UP
CALL TO ORDER—9:00 am	
Introductions	
Public Comment – None	
APPROVAL OF MINUTES – S. Lieberman motioned, R. Colombo 2nd. Approved.	
DISCUSSION ITEMS	
Addition of Ketamine <ul style="list-style-type: none"> - In the effort to expand the current toolbox of pain medications for our county's patients, efforts were made to investigate and develop a LOSOP for Ketamine. - SLOEMSA is wanting to renew efforts to add Ketamine to SLOEMSA's protocols and formulary. - Ketamine has successfully passed in Clinical Advisory Committee. - Following recommendation in Operations, Ketamine would be sent to EMCC for review and implementation in July 1st, 2024. - Further amendments: <ul style="list-style-type: none"> o Change formatting, add an "or" to the wording. o Note: could start with Ketamine and if it is a long transport could follow up with Fentanyl. o Be specific on the size of the vial. o Add an option for an additional dose. 	R. Rosander
Discussion R. Jenkins and K. Strommen confirm that there are spots in the kits to add Ketamine. R. Colombo asks why some large counties are holding off on using Ketamine. R. Colombo asks if they can transfer the controlled substance without documentation or with minimal documentation. R. Jenkins says that we need to be very clear with Paramedics that this is for pain management, not behavioral. R. Jenkins makes a motion to approve with changes.	Motion to approve: R. Jenkins 2 nd : K. Strommen Approved
Items Moving Forward Tactical EMS and Scene Management Public Safety First Aid	
ADJOURN – 9:31 am	
Next Meeting: April 4th, 2024, 09:00 A.M. Location: SLO EMSA - 2995 McMillan Ave, Suite 178, San Luis Obispo	

AGENDA ITEM / DISCUSSION	ACTION / FOLLOW-UP



COUNTY OF SAN LUIS OBISPO HEALTH AGENCY

PUBLIC HEALTH DEPARTMENT

Penny Borenstein, MD, MPH Health Officer/Public Health Director

MEETING DATE	April 4 th , 2024
STAFF CONTACT	Ryan Rosander, EMS Coordinator 805.788.2513 rrosander@co.slo.ca.us
SUBJECT	Airway/Cardiac Arrest Management
SUMMARY	<p>After implementing SGA in the County of San Luis Obispo on 07/01/2023, discussions with multiple stakeholders have occurred about the confusion surrounding when to initiate a supraglottic airway (SGA), especially for cardiac arrest. Furthermore, SGA adoption went through the committee process as a perceived backup airway to endotracheal intubation (ETI). After multiple discussions, SLOEMSA has decided to send SGA, ETI, airway management, and atraumatic/traumatic cardiac arrest management back through the committee process for further clarification. The changes are as follows;</p> <p>Protocol #602: Airway Management</p> <ul style="list-style-type: none"> • Adding provider discretion for which ALS airway to use, ETI or SGA. • Removed all language about first visualizing a patient’s airway/vocal cords before determining which ALS airway to utilize. <p>Procedure #717: Endotracheal Intubation</p> <ul style="list-style-type: none"> • Revised ETI indications to include cardiac arrest regardless of ROSC. • Removed situations where airway cannot be maintained by BLS techniques from indications list. • Removed language about BLS airway use, this is covered in BLS protocols. • Added after 2nd ETI attempt the provider shall proceed to SGA. • Added the definition of compromised airway in reference to ETI indications. • Added ETI is indicated during cardiac arrest if provider feels they can do so without interruption in HPCPR otherwise, proceed directly to SGA. • Added PCR documentation component if ALS airway cannot be established. <p>Procedure #718: Supraglottic Airway Device</p> <ul style="list-style-type: none"> • Removed all language about having to first visualize a patient’s airway/vocal cords before SGA utilization. • Added SGA is indicated in cardiac arrest. • Added PCR documentation component if ALS airway cannot be established. • Removed all language about having to first visualize a patient’s airway/vocal cords and then determine which ALS airway to utilize.

Emergency Medical Services

2995 McMillan Way Suite 178 | San Luis Obispo, CA 93401 | (P) 805-781-2519 | (F) 805-788-2517

www.slocounty.ca.gov/ems

	<p>Protocol #641: Cardiac Arrest (Atraumatic)</p> <ul style="list-style-type: none"> • Adding provider discretion to ETI or SGA utilization but shall utilize ALS airway. • Removing ROSC language to ALS airway utilization. • Added PCR documentation component if ALS airway cannot be established. <p>Protocol #661 Traumatic Cardiac Arrest</p> <ul style="list-style-type: none"> • Shall utilize Oral Intubation or Supraglottic Airways (Adults), provider discretion. • Added PCR documentation component if ALS airway cannot be established. <p>Following approval, revisions to protocols #602, #641, #661 and procedures #717 and #718 would be sent to EMCC for Adoption. Potential implementation date would be after training occurring during the 2024 SLOEMSA Update Class.</p>
REVIEWED BY	Dr. William Mulkerin, SLOEMSA Staff, Clinical Advisory Committee
RECOMMENDED ACTION(S)	Recommended revisions to protocols #602, #641, #661 and procedures #717, #718 for adoption by Operations and move to EMCC Agenda.
ATTACHMENT(S)	Protocols: #602, #641, #661 Procedures: #717, #718

AIRWAY MANAGEMENT	
ADULT	PEDIATRIC (<34 kg)
BLS	
<ul style="list-style-type: none"> • Universal Protocol #601 • Administer O₂ as clinical symptoms indicate (see notes below) • Pulse oximetry • Patients with O₂ Sat ≥ 94% without signs or symptoms of hypoxia or respiratory compromise should not receive O₂ • When applying O₂ use the simplest method to maintain O₂ Sat ≥ 94% • Do not withhold O₂ if patient is in respiratory distress • Foreign Body/Airway Obstruction <ul style="list-style-type: none"> ○ Use current BLS choking procedures ○ Basic airway adjuncts and suctioning as indicated and tolerated 	<p style="text-align: center;">Same as Adult (except for newborns)</p> <ul style="list-style-type: none"> • Newborn (< 1 day) follow AHA guidelines – Newborn Protocol #651
BLS Elective Skills	
<ul style="list-style-type: none"> • Moderate to Severe Respiratory Distress <ul style="list-style-type: none"> ○ CPAP as needed – CPAP procedure #703 	<p style="text-align: center;">CPAP not used for patients ≤34 kg</p>
ALS Standing Orders	
<ul style="list-style-type: none"> • Foreign Body/Airway Obstruction If obstruction not relieved with BLS maneuvers <ul style="list-style-type: none"> ○ Visualize and remove obstruction with Magill forceps ○ If obstruction persists, consider – Needle Cricothyrotomy Procedure #704 ○ Upon securing airway monitor O₂ Sat and ETCO₂ – Capnography Procedure #701 • Endotracheal intubation – as indicated to control airway – Procedure #717 • Supraglottic Airway – as indicated to control airway if indicated – Procedure #718 • Needle thoracostomy with symptoms of tension pneumothorax – Needle Thoracostomy Procedure #705 	<ul style="list-style-type: none"> • Foreign Body/Airway Obstruction If obstruction not relieved with BLS maneuvers <ul style="list-style-type: none"> ○ Visualize and remove obstruction with Magill forceps ○ If obstruction persists, consider – Needle Cricothyrotomy Procedure #704 ○ Upon securing airway monitor O₂ Sat and ETCO₂ – Capnography Procedure #701 • Needle thoracostomy with symptoms of tension pneumothorax – Needle Thoracostomy Procedure #705
Base Hospital Orders Only	
<ul style="list-style-type: none"> • Symptomatic Esophageal Obstruction <ul style="list-style-type: none"> ○ Glucagon 1mg IV followed by rapid flush. Give oral <u>fluid</u> challenge 60 sec after admin - check a blood sugar prior • As needed 	<ul style="list-style-type: none"> • Symptomatic Esophageal Obstruction <ul style="list-style-type: none"> ○ Glucagon 0.1mg/kg IV not to exceed 1mg followed by rapid flush. Give oral <u>fluid</u> challenge 60 sec after admin - check a blood sugar prior

	<ul style="list-style-type: none"> As needed
Notes	
<ul style="list-style-type: none"> Oxygen Delivery <ul style="list-style-type: none"> Mild distress – 0.5-6 L/min nasal cannula Severe respiratory distress – 15 L/min via non-rebreather mask Moderate to severe distress – CPAP 3-15 cm H2O Assisted respirations with BVM – 15 L/min Pediatric intubation is no longer an approved ALS skill – maintain with BLS options Patients requiring an advanced airway, providers shall decide which ALS airway to utilize based on discretion the complexity of the patient's anatomy. If the patient's vocal cords are easily visualized, then Endotracheal Intubation shall be utilized. If the patient's vocal cords are difficult or unable to be visualized, then a Supraglottic Airway Device shall be utilized. During assessments of an airway for advanced airway placement, an attempt at visualization shall be defined as placement of a laryngoscope blade and the lifting of the patient's jaw in order to visualize vocal cords. An attempt at ETI shall be defined as attempting to pass the tube through the patient's vocal cords without success. After placement of any advanced airway, providers shall verify placement of the advanced airway by waveform capnography and a minimum of one additional method. This additional method can be any of the following: <ul style="list-style-type: none"> Auscultation of lung and stomach sounds. Colorimetric CO2 Detector Device. Esophageal Bulb Detection Device. 	

Endotracheal Intubation**FOR USE IN PATIENTS >34 KG****BLS**

Universal Protocol #601

Pulse Oximetry – O₂ administration per Airway Management Protocol #602**ALS Standing Orders**

- Indications:
 - Patients with a respiratory compromise.
 - ~~ROSC~~-Patients requiring airway stabilization, including cardiac arrest and ROSC.
 - ~~Situations where the airway cannot be adequately maintained by BLS techniques.~~
- Contraindications:
 - Intact gag reflex
 - ~~If patient presents with an easily accessible airway (able to visualize the patient's vocal cords), ETI will be indicated.~~
- Prepare, position, and oxygenate the patient with 100% Oxygen. Ideal positioning is keeping the ears in line with the sternal notch.
- Consider use of video laryngoscopy when available.
- Select appropriate size ET tube and consider the need for endotracheal introducer (Bougie); have suction ready.
- Using the laryngoscope, visualize vocal cords.
- Determine how accessible the patient's airway is. If the patient has a complex airway (unable to visualize the vocal cords due to surrounding anatomy) which would be difficult and time consuming to intubate, consider the use of a supraglottic airway device Procedure # 718.
- Visualization of vocal cords will take no longer than 10 seconds.
- Visualize tube/bougie passing through vocal cords.
- Inflate the cuff with 3-10mL of air.
- Apply waveform capnography (reference Policy #701).
- Auscultate for bilaterally equal breath sounds and absence of sounds over the epigastrium.
- If ET intubation efforts are unsuccessful after the 1st attempt, ~~continue with a BLS airway,~~ oxygenate and re-evaluate the airway positioning before the 2nd attempt. After first failed attempt, consider use of Supraglottic Airways (reference Procedure #718).
- If ET intubation efforts are unsuccessful after the 2nd attempt, oxygenate and ~~continue with a BLS airway and~~ provider shall then proceed to Supraglottic Airway Procedure #718.
- Patients who have an advanced airway established shall have that airway secured with tape or a commercial device. Devices and tape should be applied in a manner that avoids compression of the front and sides of the neck, which may impair venous return from the brain.

- If the patient has a suspected spinal injury:
 - Open the airway using a jaw-thrust without head extension.
 - If airway cannot be maintained with jaw thrust, use a head-tilt/chin-lift maneuver.
 - Manually stabilize the head and neck rather than using an immobilization device during CPR.
- Following placement of the Endotracheal Tube, if the patient is noted to have an ETCO₂ less than 10, the ALS Provider shall extubate the patient and oxygenate prior to an additional attempt.

Base Hospital Orders Only

As needed

Notes

- Respiratory compromise is defined as any condition that prevents the movement of oxygenated air into and out of the lungs. This includes cardiac arrests
- ETI during cardiac arrest is indicated if the ALS provider can accomplish intubation without interruption in HPCPR. With ALS provider judgement, determines ETI cannot be accomplished, provider shall proceed to Supraglottic Airway Procedure #718
- Once an SGA has been placed, it should not be removed for an ETI
- If the provider cannot accomplish an ALS airway, they should document in the PCR why an ALS airway wasn't accomplished
- ~~During the initial visualization of the patient's airway if the ALS provider determines the airway to be difficult (unable to visualize the patient's vocal cords), ETI will not be utilized and ALS providers will reference Procedure 718 for SGA.~~
- After placement of the Endotracheal Tube, providers shall verify placement of the ETI by waveform capnography and a minimum of one additional method. This additional method can be any of the following:
 - Auscultation of lung and stomach sounds.
 - Colorimetric CO₂ Detector Device
 - Esophageal Bulb Detection Device
 - During placement of an ETI, apneic oxygenation is recommended to be utilized when available. If appropriate, providers shall place a nasal cannula onto the patient prior to the intubation attempt and continue use of the nasal cannula during placement to assist in oxygenation

Supraglottic Airway Device															
FOR USE IN PATIENTS >34 KG															
BLS															
Universal Protocol #601															
Pulse Oximetry – O ₂ administration per Airway Management Protocol #602															
ALS Standing Orders															
<ul style="list-style-type: none"> Patients who meet indications for Endotracheal Intubation Procedure #717 Patients who after the ALS Provider has visualized the patient's airway and has determined that their airway will be difficult to access. ALS provider judgement. SGA use is not approved for pediatric use. SGA shall only be used for patients >34kg. 															
I-GEL															
<ul style="list-style-type: none"> Monitor End-tidal capnography throughout use. Select appropriate tube size. <table border="1" style="margin-left: auto; margin-right: auto;"> <tbody> <tr> <td style="background-color: yellow; width: 30px;"></td> <td style="text-align: center;">3</td> <td style="text-align: center;">Small Adult</td> <td style="text-align: center;">30-60kg</td> </tr> <tr> <td style="background-color: green; width: 30px;"></td> <td style="text-align: center;">4</td> <td style="text-align: center;">Medium Adult</td> <td style="text-align: center;">50-90kg</td> </tr> <tr> <td style="background-color: orange; width: 30px;"></td> <td style="text-align: center;">5</td> <td style="text-align: center;">Large Adult</td> <td style="text-align: center;">90+kg</td> </tr> </tbody> </table> <ul style="list-style-type: none"> While preparing tube, have assistive personnel open the airway, and clear of any foreign objects. Pre-oxygenate with 100% oxygen via bls airway and BVM. Apply water soluble lubricant to the distal tip and posterior aspect (only) of the tube, taking care to avoid introduction of the lubricant into or near the ventilatory openings. Grasp the lubricated i-gel firmly along the integral bite block. Position the device so that the i-gel cuff outlet is facing towards the chin of the patient. Position patient into "sniffing position" with head extended and neck flexed. The chin should be gently pressed down before proceeding to insert the i-Gel. Introduce the leading soft tip into the mouth of the patient in the direction towards the hard palate. Glide the device downwards and backwards along the hard palate with a continuous but gentle push until a definitive resistance is felt. At this point the tip of the airway should be located into the upper esophageal opening and the cuff should be located against the laryngeal framework. The incisors should be resting on the integral bite-block. Attach a BVM. While gently bagging the patient to assess ventilation, carefully withdraw the airway until ventilation is easy and free flowing (large tidal volume with minimal airway pressure). Confirm proper position by auscultation, chest movement and verification of ETCO₂ by waveform capnography. The i-gel should be secured down per manufacturer recommendation. Patients who have an advanced airway established shall have that airway secured with tape or a commercial device. Devices and tape should be applied in a manner that avoids compression of the front and sides of the neck, which may impair venous return from the brain. 					3	Small Adult	30-60kg		4	Medium Adult	50-90kg		5	Large Adult	90+kg
	3	Small Adult	30-60kg												
	4	Medium Adult	50-90kg												
	5	Large Adult	90+kg												

- Ensure proper documentation of placement of the i-Gel placement including verification methods.

Base Hospital Orders Only

As needed

Notes

Contraindications

- Gag reflex. •Caustic ingestion. •Known esophageal disease (e.g., cancer, varices, or stricture).

- SGA during cardiac arrest is indicated
- Once an SGA has been placed, it should not be removed for an ETI
- If the provider cannot accomplish an ALS airway, they should document in the PCR why an ALS airway wasn't accomplished
- ~~Following visualization of the patient's airway and determining the patient's airway to be accessible (able to visualize the patient's vocal cords), SGA shall not be utilized and ALS providers shall reference Procedure #717 for ETI.~~
- To verify patency and placement of the SGA Device, providers shall verify placement of the i-Gel device by waveform capnography and a minimum of one additional method. This additional method can be any of the following:
 - Auscultation of lung sounds
 - Colorimetric CO2 Detector Device
 - Esophageal Bulb Detection Device
- During placement of an SGA, apneic oxygenation is recommended to be utilized when available. If appropriate, providers shall place a nasal cannula onto the patient prior to i-Gel placement and continue use of the nasal cannula during placement in order to assist in oxygenation.

CARDIAC ARREST (ATRAUMATIC)	
ADULT	PEDIATRIC (≤34 KG)
BLS Procedures	
<ul style="list-style-type: none"> • Universal Algorithm #601 • High Performance CPR (HPCPR) (10:1) per Procedure #712 <ul style="list-style-type: none"> • Continuous compressions with 1 short breath every 10 compressions • AED application (if shock advised, administer 30 compressions prior to shocking) • Pulse Oximetry <ul style="list-style-type: none"> • O₂ administration per Airway Management Protocol #602 	<ul style="list-style-type: none"> • Same as Adult (except for neonate) • Neonate (<1 month) follow AHA guidelines • CPR compression to ventilation ratio <ul style="list-style-type: none"> • Newborn – CPR 3:1 • 1 day to 1 month – CPR 15:2 • >1 month – HPCPR 10:1 • AED – pediatric patient >1 year • Use Broselow tape or equivalent if available
ALS Procedures	
<p style="text-align: center;">Rhythm analysis and shocks</p> <ul style="list-style-type: none"> • At 200 compressions begin charging the defibrillator while continuing CPR • Once fully charged, stop CPR for rhythm analysis • Defibrillate V-Fib/Pulseless V-tach – Shock at 120J and immediately resume CPR <ul style="list-style-type: none"> • Subsequent shock, after 2 mins of CPR: 150J, then 200J • Recurrent V-fib/Pulseless V-tach use last successful shock level • No shock indicated – dump the charge and immediately resume CPR <p style="text-align: center;">V-Fib/Pulseless V-Tach and Non-shockable Rhythms</p> <ul style="list-style-type: none"> • Epinephrine 1:10,000 1mg IV/IO repeat every 3-5 min <ul style="list-style-type: none"> • Do not give epinephrine during first cycle of CPR <p style="text-align: center;">V-Fib/Pulseless V-Tach</p> <ul style="list-style-type: none"> • Lidocaine 1.5mg/kg IV/IO repeat once in 3-5 min (max total dose 3 mg/kg) 	<ul style="list-style-type: none"> • <u>Emphasize resuscitation and HPCPR rather than immediate transport</u> <p style="text-align: center;">Rhythm analysis and shocks</p> <ul style="list-style-type: none"> • Coordinate compressions and charging same as adult • Defibrillate V-Fib/Pulseless V-Tach – shock at 2 J/kg and immediately resume CPR <ul style="list-style-type: none"> • Subsequent shock, after 2 mins of CPR: 4J/kg • Recurrent V-Fib/Pulseless V-tach use last successful shock level • No shock indicated – dump the charge and immediately resume CPR <p style="text-align: center;">V-Fib/Pulseless V-Tach and Non-shockable Rhythms</p> <ul style="list-style-type: none"> • Epinephrine 1:10,000 0.01 mg/kg (0.1 ml/kg) IV/IO not to exceed 0.3mg, repeat every 3-5 min <ul style="list-style-type: none"> • Do not give epinephrine during first cycle of CPR <p style="text-align: center;">V-Fib/Pulseless V-Tach</p> <ul style="list-style-type: none"> • Lidocaine 1 mg/kg IV/IO repeat every 5 min (max total dose 3 mg/kg)
Base Hospital Orders Only	
<p style="text-align: center;">ROSC with Persistent Hypotension</p> <ul style="list-style-type: none"> • Push-Dose Epinephrine 10 mcg/ml 1ml IV/IO every 1-3 min 	<p>Contact closest Base Hospital for additional orders</p> <p style="text-align: center;">ROSC with Persistent Hypotension for Age</p>

<ul style="list-style-type: none"> Repeat as needed titrated to SBP >90mmHg <u>See notes for mixing instructions</u> <p style="text-align: center;"><u>OR</u></p> <ul style="list-style-type: none"> Epinephrine Drip start at 10 mcg/min IV/IO infusion <ul style="list-style-type: none"> Consider for extended transport <u>See formulary for mixing instructions</u> <p>Contact STEMI Receiving Center (French Hospital)</p> <ul style="list-style-type: none"> Refractory V-Fib or V-Tach not responsive to treatment Request for a change in destination if patient rearrests en route Termination orders when unresponsive to resuscitative measures As needed <p>Contact appropriate Base Station per Base Station Report Policy #121 – Atraumatic cardiac arrests due to non-cardiac origin (OD), drowning, etc.)</p>	<ul style="list-style-type: none"> Push-Dose Epinephrine 10 mcg/ml 1 ml IV/IO (0.1 ml/kg if <10kg) every 1-3 min <ul style="list-style-type: none"> Repeat as needed titrated to age appropriate SBP <u>See notes for mixing instructions</u> <p style="text-align: center;"><u>OR</u></p> <ul style="list-style-type: none"> Epinephrine Drip start at 1 mcg/min, up to max of 10 mcg/min IV/IO infusion <ul style="list-style-type: none"> Consider for extended transport <u>See formulary for mixing instructions</u> As needed
---	--

Notes
<ul style="list-style-type: none"> Mixing Push-Dose Epinephrine 10 mcg/ml (1:100,000): Mix 9 ml of Normal Saline with 1 ml of Epinephrine 1:10,000, mix well. Use manufacturer recommended energy settings if different from listed Assess for reversible causes: tension PTX, hypoxia, hypovolemia, hypothermia, hyperkalemia, hypoglycemia, overdose Vascular access – IV preferred over IO – continue vascular access attempts even if IO access established) Shall utilize Oral Intubation or Supraglottic Airways (Adults), provider discretion —Utilize if airway is not patent or with maintained ROSC During the initial visualization of the patient’s airway if If the provider cannot accomplish an ALS airway, they should document in the PCR why an ALS airway wasn’t accomplished Once an SGA has been placed, it should not be removed for an ETI Stay on scene to establish vascular access, provide for airway management, and administer the first dose of epinephrine followed by 2 min of HPCPR Adult ROSC that is maintained: <ul style="list-style-type: none"> Obtain 12-lead ECG and vital signs Transport to the nearest STEMI Receiving Center regardless of 12-lead ECG reading Maintain O2 Sat greater than or equal to 94% Monitor ETCO2 Protect airway with oral intubation or Supraglottic Airway

- With BP < 100 mmHg, contact SRC (French Hospital) for fluid, or pressors
- Termination for patients > 34 kg – Contact SRC (French Hospital) for termination orders
- If the patient remains pulseless and apneic following 20 minutes of resuscitative measures
- Persistent ETCO₂ values < 10 mmHg, consider termination of resuscitation
- Documentation shall include the patient's failure to respond to treatment and of a non-viable cardiac rhythm (copy of rhythm strip)
- Pediatric patients less than or equal to 34 kg
- Evaluate and treat for respiratory causes
- Use Broselow tape if available
- Contact and transport to the nearest Base Hospital
- Receiving Hospital shall provide medical direction/termination for pediatric patients

DRAFT

TRAUMATIC CARDIAC ARREST	
ADULT	PEDIATRIC (≤34KG)
BLS	
<ul style="list-style-type: none"> • Universal Protocol #601 • Obvious Death – see Prehospital Determination of Death Policy #125 • Follow HPCPR guidelines for CPR (10:1) and minimize interruptions (< 5 seconds) 	Same as Adult
BLS Optional	
Pulse Oximetry – O ₂ administration per Airway Management Protocol #602	
ALS Standing Orders	
<p>Traumatic arrest <u>with</u> signs of life on EMS arrival and < 20 min from trauma center or hospital</p> <ul style="list-style-type: none"> • Do not delay transport • Perform ALS treatments en route • Normal Saline up to 500 mL – repeat x1 if no ROSC or SBP of < 90 mmHg • Do not use Epinephrine or Lidocaine unless the arrest is suspected to be of medical origin • Resuscitate and treat for reversible causes, i.e. hypoxia, hypovolemia, tension pneumothorax • For suspected tension pneumothorax see Needle Thoracostomy Procedure #705 <p style="text-align: center;">Traumatic arrest <u>with absent</u> signs of life on EMS arrival</p> <ul style="list-style-type: none"> • With absent signs of life consider non-initiation – Prehospital Determination of Death Policy #125 	<p style="text-align: center;">Same as Adult (except as noted below)</p> <ul style="list-style-type: none"> • Normal Saline 20 mL/kg IV/IO – reassess and repeat
Base Hospital Orders Only	
<ul style="list-style-type: none"> • Traumatic arrest <u>with</u> signs of life on EMS arrival and > 20 min from trauma center or hospital <ul style="list-style-type: none"> ○ Contact SLO Trauma Center for treatment and/or destination • Termination of resuscitation • As needed 	Same as Adult
Notes	
<ul style="list-style-type: none"> • Absent signs of life assessment include: pulseless, apneic, lack of heart and lung sounds, fixed and dilated pupils • Trauma Center is the preferred destination if equal or near equal distance • Do not delay transport for advanced airway or other treatment modalities • Consider medical origin in older patients with low probable mechanism of injury • Unsafe scene or other circumstances may warrant transport despite low potential for survival • Minimize disturbance of potential crime scene • Shall utilize Oral Intubation or Supraglottic Airways (Adults), provider discretion 	

- If the provider cannot accomplish an ALS airway, they should document in the PCR why an ALS airway wasn't accomplished