

<b>ABDOMINAL/FLANK PAIN (MEDICAL)</b>	
<b>ADULT</b>	<b>PEDIATRIC (≤34 KG)</b>
<b>BLS</b>	
<ul style="list-style-type: none"> <li>• Universal Protocol #601</li> <li>• Pulse Oximetry                             <ul style="list-style-type: none"> <li>○ O<sub>2</sub> administration per Airway Management Protocol #602</li> </ul> </li> <li>• NPO and position of comfort</li> <li>• Monitor for orthostatic changes</li> <li>• If VS are unstable                             <ul style="list-style-type: none"> <li>○ Place supine as tolerated</li> </ul> </li> </ul>	<p>Same as Adult</p>
<b>ALS Standing Orders</b>	
<ul style="list-style-type: none"> <li>• <b>Hypotension or other signs of early shock</b> – Shock (Medical) Protocol #619</li> <li>• <b>Moderate or Severe Pain</b> – Pain Management Protocol #603</li> <li>• <b>Severe Nausea</b> – Severe Nausea/Vomiting Protocol #615</li> <li>• Consider obtaining 12-lead and blood glucose</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Evaluate signs of hypotension specific to age</b> – see Universal Protocol #601 Attachment A</li> <li>• <b>Obtain blood glucose</b></li> <li>• Same as Adult</li> </ul>
<b>Base Hospital Orders Only</b>	
<ul style="list-style-type: none"> <li>• As needed</li> </ul>	<ul style="list-style-type: none"> <li>• As needed</li> </ul>
<b>Notes</b>	
<ul style="list-style-type: none"> <li>• Consider acute coronary syndrome with atypical presentation</li> </ul>	